

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

Shao Phan

OFFICE OF CHEMICAL SAFETY
AND POLLUTION
PREVENTION

November 07, 2019

MEMORANDUM

Subject: Efficacy Review for Concrobium Mold Control; EPA File No. 82552-G; DP Barcode: D454099;

Submission #: 1036690; E-Sub # 40557.

From: Ibrahim Laniyan, Ph.D.

Microbiologist

Product Science Branch

Antimicrobials Division (7510P)

Thru: Thao Pham, Acting Efficacy Team Leader

Product Science Branch

Antimicrobials Division (7510P)

Date signed: 11/20/19

To: Jacqueline Hardy RM 34/ Stacey Grigsby

Regulatory Management Branch II Antimicrobials Division (7510P)

Applicant: Siamons International, Inc.

48 Galaxy Blvd., Unit 413 Toronto. Ontario M9W 6C8

Canada

Formulation from the Label:

Active Ingredients	% by wt.
Sodium Carbonate	1.005 %
Other Ingredients:	98.995 %
Total	100.000 %

I. BACKGROUND

Product Description (as packaged, as applied): Liquid RTU.

Submission type: New end use product

Currently registered efficacy claim(s): NA

Requested action(s): New product Registration

Documents considered in this review:

• Letter from applicant to EPA dated June 26, 2019

- Application for Pesticide (EPA form 8570-1) dated June 26, 2019
- Confidential Statement of Formula (EPA form 8570-4) dated June 26, 2019
- Certification with Respect to Citation of Data (EPA Form 8570-34) dated June 26, 2019
- Data Matrix (EPA Form 8570-35) dated June 26, 2019
- 5 efficacy studies (MRID nos. 501633-01 501633-09)
- Proposed label dated 05/26/2019.

II. PROPOSED DIRECTIONS FOR USE

[[Remove heavy soil prior to application] [Mold stains may remain visible following treatment] [This is normal] [In this case] [If so] [If mold stains remain] [following treatment] [reapply] [product] [and] [wipe] [or] [scrub] [area] [away stains] [In some cases, staining may be irreversible.] [[Do not expect] [Concrobium Mold Control [insert brand name] [this product] [to] [will not] [whiten] stains [on contact]]. [If, [following treatment], additional steps are taken to [remove] [or] [cover] [address] [treat] the [mold] stain, reapply [[Concrobium Mold Control [insert brand name] [this product]] as a final step [as] [per] [as indicated in] [the] [Preventative Treatment] [To Prevent] [of] [[mold] [and] [mildew] section] [to protect area from regrowth.]]

Apply [Concrobium Mold Control [insert brand name] [this product]] by cloth[,] [or] [sponge,], handspray, [electric or compression sprayer,] [paintbrush,] [roller,] [or immersion]. Expect 350-2800 sq. ft. of surface coverage per gallon depending on surface type [and application method].

[FOR CLEANING [AND] [DEGREASING]] {numbering of steps is optional}

- [1)] Apply [product] [Concrobium Mold Control [insert brand name]] on area until wet.
- [2)] Wipe [or scrub][Wipe/scrub] clean [and dry]. [No rinsing required.]
- [3)] [For tough stains, reapplication may be necessary.]]

[PREVENTATIVE TREATMENT] [FOR] [TO PREVENT] [OF] [MOLD] [AND] [MILDEW]

To [inhibit] [prevent] mold and mildew growth on surfaces:

{numbering of steps is optional}

- [1)] Apply [product] [Concrobium Mold Control [insert brand name]] on area until evenly wet.
- [2)] [[If desired][If dry time][If water staining][is a] [potential] [concern] [To promote drying] [and] [to][avoid residue marks], [use a] [clean cloth] [or] [paper towel] [dampened with [product] Concrobium Mold Control [insert brand name]] [to] [wipe] [off] [wipe off] [clean] [clean off] [off excess wetness] [with a clean cloth or paper towel].
- [3)] Allow treated surfaces to dry completely. [The product generally dries within 2 hours and may take up to 24 hours depending on surface absorbency, temperature and [drying] conditions.]
- [[4)] [If treated surfaces are to be [cleaned] [and] [/] [or] [painted], allow [product] [Concrobium Mold Control [insert brand name]] to dry for 24 hours.]

To Disinfect [and] [Kill Fungi [& Mildew]]: {numbering of steps is optional}

- [1] Clean surfaces to remove visible soil.
- [2] From a distance of 6-8", spray product on hard, nonporous surfaces until thoroughly wet. Allow to remain wet for 10 minutes.
- [3] [Wipes surfaces] [and] [or] [allow to air dry].

III. STUDY SUMMARIES

1.	MRID	508217-11	508217-11 Study Completion Date: October 25, 2018				
Study (Objective	Disinfectant – Bactericidal					
Testing Lab	, Lab Study ID		Accuratus	Lab Serv	rices, A25891		
Test org	ganism(s)	Pseudomonas a	eruginosa (ATC	C 15442),	Salmonella ent	erica (A	TCC 10708),
□ 1 □ 2 ⊠ 3 □ 4+ and Staphylococcus aureus (ATCC 6538)					88)		
Test	Method	AOAC Germicida	. ,	copy provi	ided)		
Applicati	on Method	Carriers were individually sprayed 15 times (15 sprays) or until thoroughly wet 30 centimeters				roughly wet at	
	Name/ID		CMC-Plus (Concrobium Mold Control)				
Test Substance	Lots □ 1 □ 2 ⊠ 3	Lot# SII-23022018-A, Lot# SII-23022018-B, and Lot# SII-23022018-C					
Preparation	Preparation	Conveniently diluted to the LCL with sterile deionized water				ater	
Soi	l load		No	soil load re	equired		
Carrier ty	pe, # per lot	Glass slides, 60 per microorganism and per batch					
				RH	46-55%		
Neut	ralizer	Letheen Broth + 0.07% Lecithin + 0.5% Tween 80 + 0.1% Sodium Thiosulfate (20 ml)					
(i.e. protocol amendments,	comments deviations and retesting, control utralizer, etc.)					8-36.0°C) and dry. pelow the osa.	

2.	MRID	508217-12	Study Completion Date:	April 15, 2019		
Study (Objective	Disinfectant – Fungicide				
Testing Lab, Lab Study ID Microchem, GLP2071						
Test organism(s) ⊠ 1 □ 2 □ 3 □ 4+		Trichophyton interdigitale (ATCC 9533)				
Test I	Method	AOAC Germicidal Spray Products as Disinfectants Test - Microchem Laboratory Protocol # P2309 (copy provided)				
Application Method		Carriers were individually sprayed 15 times (15 sprays) from 6-8 inches away at approximately 45° from the test carrier to the spray nozzle.				
Name/ID		CMC-Plus (Concrobium Mold Control)				
Test Substance	Lots □ 1 ⊠ 2 □ 3	SII-18122018-B, SII-18122018-C				
Preparation Preparation		Conveniently diluted to the LCL with sterile deionized water				
Soi	load	No soil load required				
Carrier ty	pe, # per lot		Glass slides, 10 per microorganism and per	batch		

NeutralizerSabouraud Dextrose Broth supplemented to contain 0.07% Lecithin and 0.5% Tween 80Carriers were each inoculated with 10.0 μL of culture and were allowed to dry for 30-31 minutes at 35-37°C until visibly dry.Reviewer comments (i.e. protocol deviations and amendments, retesting, control failures, neutralizer, etc.)Protocol was amended to add a test with an increased contact time. After failure at 9 minutes 50 seconds contact time, lot SII-18122018-C was tested using a 9 minute 55 second exposure time.	Test conditions	Contact time	9min50sec 9min55 sec.	Temp	20.4-23.4°C	RH	-
for 30-31 minutes at 35-37°C until visibly dry. Reviewer comments (i.e. protocol deviations and amendments, retesting, control at 9 minutes 50 seconds contact time, lot SII-18122018-C was tested using a 9 minute 55 second exposure time.	Neutralizer	Sabouraud Dextrose Broth supplemented to contain 0.07% Lecithin and 0.5% Tween 80					
The test substance was applied using 15 trigger sprays for testing. More	(i.e. protocol deviations and amendments, retesting, control	Protocol was ar at 9 minutes 50	for 30-31 minute mended to add a t 0 seconds contact minute 55	es at 35- est with a time, lot second	37°C until visibly an increased con SII-18122018-C exposure time.	dry. tact time was tes	e. After failure sted using a 9

3.	MRID	508217-13	Study Co	ompletio	n Date:	Apr	il 23, 2019
Study (Objective	Disinfectant – Fungicide					
Testing Lab	, Lab Study ID	Accuratus Lab Services, A25595					
	ganism(s) : □ 3 □ 4+		Aspergil	lus niger ((ATCC 6275)		
Test	Method	Fungicidal G	Sermicidal Spray N SIA0010110		Accuratus Lab S I (<i>copy provided</i>		Protocol #
Applicati	Application Method Carriers were individually sprayed 15 times (15 sprays) from 30 cen away, at in an undisturbed horizontal position.			centimeters			
_ ,	Name/ID	CMC-Plus (Concrobium Mold Control)					
Test Substance	Lots □ 1 ⊠ 2 □ 3	Lot# SII-23022018-A and Lot# SII-23022018-B					
Preparation	Preparation	Conv	eniently diluted to	the LCL	with sterile deio	nized wa	ater
Soi	load		No	soil load ı	required		
Carrier ty	pe, # per lot		Glass slides, 10 p	er microo	rganism and per	batch	
Test co	onditions	Contact time	9.50 min. 9min45 sec.	Temp	20-21°C	RH	48-51%
Neut	ralizer			-			
		Carriers were each inoculated with 10.0 µL of culture and were allowed to dry for 30-31 minutes at 36.1-36.2°C under and at a 43-51.7% relative humidity, until visibly dry.					
Reviewer comments (i.e. protocol deviations and amendments, retesting, control failures, neutralizer, etc.) Protocol was amended to add an additional test substance parameter failure at 9.5 minutes contact time, lot SII-23022018-A was tested usin minute 45 second exposure time. The test substance was applied using 15 trigger sprays for testing.					d using a 9		
			information is nee				

4.	MRID	508217-14 Study Completion Date: October 10, 2018					
Study	Objective		Hard Surface Mildew Fungistatic				
Testing Lab	, Lab Study ID		Accuratus Lab Services, A25877				
	ganism(s) 2		Aspergillus niger (ATCC 6275)				
Test	Method	EPA Hard Surfa	ice Mildew-Fungistatic Test – Accuratus La SIA001011018.MSTAT (<i>copy provide</i> d				
Applicati	ion Method		e individually sprayed 15 times (15 sprays) tertical or near vertical position to permit exc				
Test	Name/ID	CMC-Plus (Concrobium Mold Control)					
Substance Preparation	Lots □ 1 ⊠ 2 □ 3	3 Lot# SII-23022018-B and Lot# SII-23022018-C					

Preparation	Conv	Conveniently diluted to the LCL with sterile deionized water					
Soil load		No soil load required					
Carrier type, # per lot	One inch by on	One inch by one inch glazed ceramic tiles, 10 per microorganism and per batch					
Test conditions	Contact time	Contact time - Temp = RH -					
Neutralizer		-					
Reviewer comments (i.e. protocol deviations and amendments, retesting, control failures, neutralizer, etc.)	Carriers were of for 26 minuted for	Carriers were dried for 20 minutes at 35.8-36.1°C under and at a 32% rehumidity Carriers were each inoculated with 3 sprays of culture and were allowed for 26 minutes at 36.1-36.2°C under and at a 28.7% relative humidity, visibly dry. Each carrier (contaminated side up) was placed onto an individual water plate. All plates were incubated for 7 days at 29.0°C in a minimum of 9 relative humidity. The test substance was applied using 15 trigger sprays for testing. Minformation is needed to support this application.					

5.	MRID	508217-15	Study C	ompletior	Date:	Octo	ber 10, 2018
Study	Objective		Fabrio	Mildew F	ungistatic		
Testing Lab	, Lab Study ID		Accuratus	s Lab Serv	rices, A25876		
	ganism(s) 2	Aspergillus	s niger (ATCC 627	75) and <i>Pe</i>	enicillium variabi	e (ATC	C 32333)
Test	Method	Fabric Mi	ildew Fungistatic ⁻ SIA00101101		uratus Lab Serv T (<i>copy provid</i> e		otocol #
Applicati	ion Method		carriers was spra ertical or near ver				
_ ,	Name/ID	CMC-Plus (Concrobium Mold Control)					
Test Substance Preparation	Lots □ 1 ⊠ 2 □ 3	Lot# SII-23022018-B and Lot# SII-23022018-C					
Freparation	Preparation	Conv	eniently diluted to	the LCL v	vith sterile deior	nized wa	ater
Soil load				soil load r			
Carrier ty	pe, # per lot		c carriers (25 mm z/yd²) cotton musli				
Test co	onditions	Contact time	-	Temp	=	RH	-
Neutralizer Sabouraud Dextrose Broth + 0.07% Lecithin and 0.5			5% Twe	een 80			
Reviewer comments (i.e. protocol deviations and amendments, retesting, control failures, neutralizer, etc.) The carriers were dried at room temperature (22.1°C) for 30 m Each side of carrier was inoculated with 3 sprays of mixed inoculated and sprays per carrier). Each contaminated carrier was suspended in individual 250 ml bottles containing approximately 10 ml sterile deionized water (fincubated at 29.0°C. Observations were made and recorded effour weeks.			inocula 50 ml Fr ater (for	(for a total of ench Square humidity) and			

V. RESULTS

MRID	Organism	No. Exhibi	ting Growth/Total	No. Tested	Average			
(Test Date)	(Test Date) SII-23022018-A SII-23022018-B SII-23022018-C log ₁₀ /Carrier							
Diluted to the LCL and 9.5 minutes contact time – No soil load								

MRID	Organism	No. Exhibit	ting Growth/Total	No. Tested	Average
(Test Date)	Organism	SII-23022018-A	SII-23022018-B	SII-23022018-C	log ₁₀ /Carrier
	Pseudomonas aeruginosa (ATCC 15442)	0/60	0/60	0/60	5.45
508217-11	Salmonella enterica (ATCC 10708)	0/60	0/60	0/60	5.62
	Staphylococcus aureus (ATCC 6538)	0/60	0/60	0/60	5.56
	Diluted to the LCL - 9.5	minutes and 9 minutes	utes 45 seconds col	ntact time – No soil	load
508217-13	Aspergillus niger (ATCC 6275)	0/10	0/10	-	4.35 / 4.37
		SII-18122018-B	SII-18122018-C	-	
Dilu	ted to the LCL - 9 minute	s 50 seconds and 9	minutes 55 second	ds contact time – No	soil load
508217-12	Trichophyton interdigitale (ATCC 9533)	0/10	0/10	-	4.02 / 4.39

MRID # 508217-14	Carrier	Carrier Visual Evaluation umber of Controls		Evaluation -23022018-B	7 Days Evaluation Lot SII-23022018-C		
500217-14	Number	of Controls	Visual	Magnified	Visual	Magnified	
	1	Growth (80%)	0%	No growth	0%	No growth	
Aspergillus niger	2	Growth (80%)	0%	No growth	0%	No growth	
	3	Growth (80%)	0%	No growth	0%	No growth	
	4	Growth (70%)	0%	No growth	0%	No growth	
	5	Growth (80%)	0%	No growth	0%	No growth	
(ATCC 6275)	6	Growth (80%)	0%	No growth	0%	No growth	
	7	Growth (80%)	0%	No growth	0%	No growth	
	8	Growth (80%)	0%	No growth	0%	No growth	
	9	Growth (90%)	0%	No growth	0%	No growth	
	10	Growth (90%)	0%	No growth	0%	No growth	

MRID # 508217-15	Carrier Number	Visual Evaluation of Controls	7 Days Evaluation Lot SII-23022018-I			
500217-15	Number	oi Collilois	Visual	Magnified	Visual	Magnified
	1	Growth (80%)	0%	No growth	0%	No growth
Mixed 50/50 Aspergillus niger	2	Growth (80%)	0%	No growth	0%	No growth
	3	Growth (80%)	0%	No growth	0%	No growth
	4	Growth (70%)	0%	No growth	0%	No growth
	5	Growth (80%)	0%	No growth	0%	No growth
(ATCC 6275) and Penicillium variable	6	Growth (80%)	0%	No growth	0%	No growth
(ATCC 32333)	7	Growth (80%)	0%	No growth	0%	No growth
	8	Growth (80%)	0%	No growth	0%	No growth
	9	Growth (90%)	0%	No growth	0%	No growth
	10	Growth (90%)	0%	No growth	0%	No growth

MRID#	Carrier	Visual Evaluation of Controls (% Coverage)						
508217-15	Number	Day 7	Day 14	Day 21	Day 28			
Mixed 50/50 Aspergillus niger (ATCC 6275) and Penicillium variable (ATCC 32333)	1	Growth (60%)	Growth (65%)	Growth (75%)	Growth (100%)			
	2	Growth (50%)	Growth (55%)	Growth (60%)	Growth (85%)			
	3	Growth (50%)	Growth (60%)	Growth (70%)	Growth (100%)			
	4	Growth (50%)	Growth (55%)	Growth (60%)	Growth (80%)			
	5	Growth (50%)	Growth (60%)	Growth 75%)	Growth (85%)			
	6	Growth (50%)	Growth (55%)	Growth (65%)	Growth (100%)			

7	Growth (50%)	Growth (50%)	Growth (60%)	Growth (100%)
8	Growth (55%)	Growth (55%)	Growth (60%)	Growth (75%)
9	Growth (50%)	Growth (50%)	Growth (60%)	Growth (100%)
10	Growth (50%)	Growth (60%)	Growth (70%)	Growth (100%)

MRID#	Conside	Mixed 50/50 Aspergillus niger (ATCC 6275) and Penicillium variable (ATCC 32333)								
508217- 15	Carrier Number	Visual	Magnified	Visual	Magnified	Visual	Magnified	Visual	Magnified	
	1	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	2	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
Evaluation	3	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
of	4	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
treatment	5	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
Lot SII-	6	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
23022018-	7	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
В	8	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	9	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	10	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	1	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	2	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
Evaluation	3	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
of	4	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
treatment	5	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
Lot SII-	6	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
23022018- C	7	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	8	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	9	0%	No growth	0%	No growth	0%	No growth	0%	No growth	
	10	0%	No growth	0%	No growth	0%	No growth	0%	No growth	

VI. CONCLUSIONS

MRID	Claim	Surface Type	Application Method(s) and Dilution	Contact Time	Soil load	Diluent	Organism(s)	Data support tested conditions?
508217-11	Disinfectant Bactericidal	Hard, non- porous surfaces	15 Sprays LCL	10 minutes	No	No	Staphylococcus aureus (ATCC 6538) Salmonella enterica (ATCC 10708) Pseudomonas aeruginosa (ATCC 15442)	No
508217-12 508217-13	Disinfectant Fungicide	Hard, non- porous surfaces	15 Sprays LCL	10 minutes	No	No	Trichophyton interdigitale (ATCC 9533) Aspergillus niger (ATCC 6275)	No
508217-14	7days Hard Surface Mildew Fungistatic	Hard, non- porous surfaces	15 Sprays LCL	-	No	No	Aspergillus niger (ATCC 6275)	No
508217-15	Fabric Mildew Fungistatic	Fabric	15 Sprays LCL	-	No	No	Aspergillus niger (ATCC 6275) and Penicillium variable (ATCC 32333)	Yes

VII. LABEL COMMENTS

Proposed Label dated 05/26/2019

Note: Please change nominal label concentrations to match the CSF.

- 1. Spray products are usually tested using 2 to 3 sprays; spraying 15 times is not representative of how a typical user would apply the product to a hard surface. More information is needed to support this excessive number of sprays, including a description of the applicator and total volume of product released from 15 sprays. Until this information is submitted and reviewed by this agency, the following claims should be removed from the label:
 - a. The proposed label claims that the product, Concrobium Mold Control (EPA File No. 82552-G), is an effective bacterial disinfectant spray, when used undiluted, at room temperature, for 10 minutes, on precleaned hard, non-porous surfaces; these claims **are unacceptable**.
 - b. The proposed label claims that the product, Concrobium Mold Control (EPA File No. 82552-G), is an effective fungicide disinfectant spray, when used undiluted, at room temperature, for 10 minutes, on precleaned hard, non-porous surfaces; these claims **are unacceptable**.
 - c. The proposed label claims that the product, Concrobium Mold Control (EPA File No. 82552-G), is an effective mildew disinfectant spray, when used undiluted, at room temperature, for 10 minutes, on precleaned hard, non-porous surfaces; these claims **are unacceptable**.
 - d. The proposed label claims that the product, Concrobium Mold Control (EPA File No. 82552-G), is an effective mildew fungistat spray, when used undiluted, at room temperature, on cleaned hard, non-porous surfaces; these claims are unacceptable. Note, for future reference, these label claims should be clearly limited to 7 days.
- 2. The proposed label claims that the product, Concrobium Mold Control (EPA File No. 82552-G), is an effective mildew fungistat spray, when used undiluted, at room temperature, on clean fabric; these claims **are acceptable.**
- 3. The product, Concrobium Mold Control (EPA File No. 82552-G) is a spray only product. Application by mop, sponge, or cloth is not acceptable.
- 4. Make the following changes to the proposed label:
 - a. Throughout the label, remove all public health claims including: disinfection, mildewcidal, fungicidal, bactericidal, mildewcide, kill(s), eliminate(s), one-step.
 - b. Remove all claims for treatment of mold including: control, inhibit, prevent, fight, defend, eradicate, abolish, destroy, crush, attack, get rid of, remediate, lethal for mold, conquer, encapsulates, tough on mold, fungal spores, etc.
 - c. Remove all claims for hard nonporous surfaces or multiple surface types. The only acceptable claim is to control mildew on clean fabric.
 - d. Remove all "2 in 1", "2 for 1", "3 in 1", "3 for 1" and "all in one" claims.
 - e. Product was tested as a spray. Remove directions for use for other application types.
 - f. Remove mold remediation language.
 - g. Revise use directions to remove public health claims and specify treatment as a mildewstat for fabrics.